

Nuclear Operators on Spaces of Continuous Vector-Valued Functions

by

Paulette Saab* and Brenda Smith

Abstract Let Ω be a compact Hausdorff space, let E be a Banach space, and let $C(\Omega, E)$ stand for the Banach space of all E -valued continuous functions on Ω under supnorm. In this paper we study when nuclear operators on $C(\Omega, E)$ spaces can be completely characterized in terms of properties of their representing vector measures. We also show that if F is a Banach space and if $T : C(\Omega, E) \rightarrow F$ is a nuclear operator, then T induces a bounded linear operator $T^\#$ from the space $C(\Omega)$ of scalar valued continuous functions on Ω into $\mathcal{N}(E, F)$ the space of nuclear operators from E to F , in this case we show that E^* has the Radon-Nikodym property if and only if $T^\#$ is nuclear whenever T is nuclear.

AMS(MOS) Subject Classification (1980)
Primary 46E40, 46G10, 47B10, Secondary 28B05, 28B20

* Research supported in part by an NSF Grant DMS-87500750

Let Ω be a compact Hausdorff space, let E be a Banach space, and let $C(\Omega, E)$ stand for the Banach space of continuous E -valued functions on Ω under supnorm. It is well known [3, p.182] that if F is a Banach space, then any bounded linear operator $T : C(\Omega, E) \longrightarrow F$ has a finitely additive vector measure G defined on the σ -field of Borel subsets of Ω with values in the space $\mathcal{L}(E, F^{**})$ of bounded linear operators from E to the second dual F^{**} of F . The measure G is said to represent T . The purpose of this note is to study the interplay between certain properties of the operator T and properties of the representing measure G . Precisely, one of our goals is to study when one can characterize nuclear operators in terms of their representing measures. This is of course motivated by a well known Theorem of L. Schwartz [5] (see also [3, p.173]) concerning nuclear operators on spaces $C(\Omega)$ of continuous scalar-valued functions. The study of nuclear operators on spaces $C(\Omega, E)$ of continuous vector-valued functions was initiated in [1] where the author extended Schwartz's result in case E^* has the Radon-Nikodym property. In this paper, we will show that the condition on E^* to have the Radon-Nikodym property is necessary to have a Schwartz's type theorem. This leads to a new characterization of dual spaces E^* with the Radon-Nikodym property. In [2] it was shown that if $T : C(\Omega, E) \rightarrow F$ is nuclear then its representing measure G takes its values in the space $\mathcal{N}(E, F)$ of nuclear operators from E to F . One of the results of this paper is that if $T : C(\Omega, E) \rightarrow F$ is nuclear then its representing measure G is countably additive and of bounded variation as a vector measure taking its values in $\mathcal{N}(E, F)$ equipped with the nuclear norm. Finally, we show by easy examples that the above mentioned conditions on the representing measure G do not characterize nuclear operators on $C(\Omega, E)$ spaces, and we also look at cases where nuclear operators are indeed characterized by the above two conditions. For all undefined notions and terminologies we refer the reader to [3].

0–Preliminaries If X and Y are Banach spaces, then $\mathcal{L}(X, Y)$ will stand for the space of bounded linear operators from X to Y . An element T in $\mathcal{L}(X, Y)$ is said to be a **nuclear operator** if there exist sequences (x_n^*) in X^* and (y_n) in Y such that for each x in X

$$T(x) = \sum_{n=1}^{\infty} x_n^*(x) y_n,$$

and

$$\sum_{n=1}^{\infty} \|x_n^*\| \|y_n\| < \infty.$$

We say that $\sum_n x_n^* \otimes y_n$ represents the nuclear operator T . The **nuclear norm** of a nuclear operator $T : X \rightarrow Y$ is defined by:

$$\|T\|_{\text{nuc}} = \inf \left\{ \sum_n \|x_n^*\| \|y_n\| \right\}$$

where the infimum is taken over all sequences (x_n^*) and (y_n) such that $T(x) = \sum_{n=1}^{\infty} x_n^*(x) y_n$ holds for all x in X . The nuclear operators from X to Y form a normed linear space under the nuclear norm [3, p.170], which we shall denote by $\mathcal{N}(X, Y)$.

If Ω is a compact Hausdorff space and E is a Banach space, then $C(\Omega, E)$ will stand for the space of continuous E -valued functions defined on Ω under supnorm. If $E = \mathbb{R}$ or $\mathbb{C}$ we will simply write $C(\Omega)$. The space $M(\Omega, E^*)$ will stand for the space of all regular E^* -valued vector measures μ defined on the σ -field Σ of Borel subsets of Ω that are of bounded variation. We shall use the fact (see [3, p.182]) that $M(\Omega, E^*)$ is a Banach space under the variation norm $\|\mu\| = |\mu|(\Omega)$, and that $M(\Omega, E^*)$ is isometrically isomorphic to the dual space $C(\Omega, E)^*$. When $E = \mathbb{R}$ or $\mathbb{C}$ we will simply write $M(\Omega)$. If $\mu \in M(\Omega, E^*)$, then for each $e \in E$ we will denote by $\langle e, \mu \rangle$ the element of $M(\Omega)$ such that for each $f \in C(\Omega)$,

$$\int f d\langle e, \mu \rangle = \mu(f \otimes e)$$

where $f \otimes e$ is the element in $C(\Omega, E)$ such that $f \otimes e(\omega) = f(\omega)e$ for each $\omega \in \Omega$.

If $\nu \in M(\Omega)$ and $x^* \in E^*$, we denote by $\nu \otimes x^*$ the element of $M(\Omega, E^*)$ which to each Borel subset B of Ω associates the element $\nu(B)x^*$ of E^* . If E and F are Banach spaces and Ω is a compact Hausdorff space, then we will denote by G the finitely additive $\mathcal{L}(E, F^{**})$ -valued measure representing the operator T . Recall that if B is a Borel subset of Ω then

$$G(B)e = T^{**}(\phi_{B,e})$$

for all $e \in E$, where $\phi_{B,e}$ is the element of $M(\Omega, E^*)^*$ such for each $\lambda \in M(\Omega, E^*)$

$$\phi_{B,e}(\lambda) = \lambda(B)(e)$$

and T^{**} is the second adjoint of T .

Finally we recall that a Banach space X has the **Radon-Nikodym property** (RNP) if for every finite measure space (S, Σ, μ) and every vector measure $m : \Sigma \rightarrow X$ of bounded variation that is absolutely continuous with respect to μ there exists a strongly measurable Bochner integrable function $g : S \rightarrow X$ such that

$$m(A) = \int_A g d\mu$$

for each $A \in \Sigma$.

1–Some Properties of the Measure representing a Nuclear operator

Throughout we let Ω be a compact Hausdorff space with Σ its σ -field of Borel subsets and we let E and F be Banach spaces. In what follows we shall look at some of the properties that a nuclear operator on $C(\Omega, E)$ induces on its representing measure G . In [2] it was shown that if $T : C(\Omega, E) \rightarrow F$ is nuclear then $G : \Sigma \rightarrow \mathcal{N}(E, F)$. In the next proposition we shall show that G enjoys a stronger property.

Proposition 1 If $T : C(\Omega, E) \rightarrow F$ is a nuclear operator with representing measure G , then:

- (i) for each Borel subset B of Ω , $G(B) : E \rightarrow F$ is a nuclear operator, and
- (ii) the measure G is countably additive and is of bounded variation as a vector measure taking its values in $\mathcal{N}(E, F)$ under the nuclear norm.

Proof: If $T : C(\Omega, E) \rightarrow F$ is nuclear, then there are sequences (μ_n) in $M(\Omega, E^*)$ and (y_n) in F such that for all $f \in C(\Omega, E)$

$$Tf = \sum_{n=1}^{\infty} \mu_n(f) y_n$$

and

$$\sum_{n=1}^{\infty} \|\mu_n\| \|y_n\| < \infty.$$

In particular the operator T^{**} is nuclear with

$$T^{**}(\phi) = \sum_{n=1}^{\infty} \phi(\mu_n) y_n$$

for all $\phi \in C(\Omega, E)^{**}$. This implies that for any Borel subset B of Ω and for all $e \in E$

$$\begin{aligned} G(B)e &= T^{**}(\phi_{B,e}) \\ &= \sum_{n=1}^{\infty} \mu_n(B)(e) y_n \end{aligned}$$

Since $(\mu_n(B))$ is a sequence in E^* and $\sum_{n=1}^{\infty} \|\mu_n(B)\| \|y_n\| \leq \sum_{n=1}^{\infty} \|\mu_n\| \|y_n\| < \infty$, this will quickly show that for each Borel subset B of Ω , the operator $G(B) : E \rightarrow F$ is nuclear. To prove (ii) note that since $G : \Sigma \rightarrow \mathcal{N}(E, F)$, let $|G|_{\text{nuc}}$ denote the extended non negative function whose value on a set B in Σ is given by

$$|G|_{\text{nuc}}(B) = \sup_{\pi} \sum_{B_i \in \pi} \|G(B_i)\|_{\text{nuc}}$$

where the supremum is taken over all finite partitions π of B . We first show that

$$|G|_{\text{nuc}}(\Omega) < \infty.$$

For this, note that if $\pi = \{B_i\}$ is a finite partition of Ω , then

$$\begin{aligned}
\sum_{B_i \in \pi} \|G(B_i)\|_{\text{nuc}} &\leq \sum_{B_i \in \pi} \sum_{n=1}^{\infty} \|\mu_n(B_i)\| \|y_n\| \\
&\leq \sum_{B_i \in \pi} \sum_{n=1}^{\infty} |\mu_n|(B_i) \|y_n\| \\
&\leq \sum_{n=1}^{\infty} \sum_{B_i \in \pi} |\mu_n|(B_i) \|y_n\| \\
&\leq \sum_{n=1}^{\infty} |\mu_n|(\Omega) \|y_n\| \\
&= \sum_{n=1}^{\infty} \|\mu_n\| \|y_n\| < \infty
\end{aligned}$$

This of course shows that $|G|_{\text{nuc}}(\Omega) < \infty$. To complete the proof of (ii) we need to show that G is countably additive. First note that since $|G|_{\text{nuc}}(\Omega) < \infty$, it follows from [3, p.7] that G is strongly additive, that is if (B_i) is a sequence of pairwise disjoint Borel subsets of Ω we have that the series $\sum_{i=1}^{\infty} G(B_i)$ converges in $\mathcal{N}(E, F)$. To complete the proof we need to check that the series $\sum_{i=1}^{\infty} G(B_i)$ converges to $G(\bigcup_{i \geq 1} B_i)$ in $\mathcal{N}(E, F)$. To this end, consider a series $\sum_{n=1}^{\infty} \mu_n \otimes y_n$ representing the operator T such that

$$\sum_{n=1}^{\infty} \|\mu_n\| \|y_n\| < \infty.$$

Without loss of generality we may and shall assume that $\|y_n\| \leq 1$ for all $n \geq 1$. Let $\epsilon > 0$ and pick $N \in \mathbb{N}$ such that

$$\sum_{\nu=N+1}^{\infty} \|\mu_n\| < \epsilon/2$$

Since $\mu_1, \mu_2, \dots$, and $\mu_N \in M(\Omega, E^*)$, there exists $K \in \mathbb{N}$ such that

$$|\mu_j| \left(\bigcup_{i \geq K} B_i \right) < \frac{\epsilon}{2^{j+1}} \text{ for all } j = 1, \dots, N$$

This implies that

$$\begin{aligned}
& \|G(\bigcup_{i \geq 1} B_i) - \sum_{i=1}^{K-1} G(B_i)\|_{\text{nuc}} = \\
& \|G\left(\bigcup_{i \geq K} B_i\right)\|_{\text{nuc}} \leq \\
& \sum_{n=1}^{\infty} \|\mu_n\left(\bigcup_{i \geq K} B_i\right)\| \leq \\
& \sum_{n=1}^N |\mu_n|\left(\bigcup_{i \geq K} B_i\right) + \sum_{n=N+1}^{\infty} |\mu_n|\left(\bigcup_{i \geq K} B_i\right) < \epsilon
\end{aligned}$$

This shows that G is countably additive as a vector measure taking its values in $\mathcal{N}(E, F)$.

The first question that arises at this stage, is when do properties (i) and (ii) above characterize nuclear operators? The next proposition shows that one necessary condition is that F should have the Radon-Nikodym property.

Proposition 2 If F fails RNP, then for any Banach space E there is a non nuclear operator $T : C([0, 1], E) \rightarrow F$ whose representing measure takes its values in $\mathcal{N}(E, F)$ and is of bounded variation as a countably additive vector measure taking its values in $\mathcal{N}(E, F)$.

Proof: If F lacks RNP then by [3, p.175] there is an F -valued countably additive vector measure m on the Borel subsets of $[0, 1]$ such that m is of bounded variation, m is absolutely continuous with respect to Lebesgue measure but m admits no Bochner integrable derivative with respect to its variation $|m|$. If we define $T' : C[0, 1] \rightarrow F$ by

$$T'(f) = \int_{[0, 1]} f dm$$

then T' is not a nuclear operator (see [5] or [3, p.173]). Now fix $e \neq 0$ in E then choose e^* in E^* with $e^*(e) = 1$ and define $T : C([0, 1], E) \rightarrow F$ by

$$T(\varphi) = T'(e^* \circ \varphi)$$

for each φ in $C([0, 1], E)$. In particular, for each f in $C[0, 1]$ and x in E we have

$$T(f \otimes x) = e^*(x)T'(f).$$

It is clear that the measure representing the operator T is the measure $G = m \otimes e^*$ which to every Borel subset B of $[0, 1]$ associates the one-rank operator such that $G(B)x = e^*(x)m(B)$ for each $x \in E$. On other hands for any finite partition $\pi = \{B_i\}$ of $[0, 1]$ we have

$$\begin{aligned} \sum_{B_i \in \pi} \|G(B_i)\|_{\text{nuc}} &\leq \sum_{B_i \in \pi} \|e^*\| \|m(B_i)\| \\ &\leq \|e^*\| |m|([0, 1]) < \infty. \end{aligned}$$

so

$$|G|_{\text{nuc}}([0, 1]) < \infty.$$

If T were a nuclear operator then there would exist (μ_n) in $M([0, 1], E^*)$ and (y_n) in F such that for each ϕ in $C([0, 1], E)$

$$T(\phi) = \sum_{n=1}^{\infty} \mu_n(\phi) y_n$$

In particular for each $f \in C[0, 1]$

$$\begin{aligned} T(f \otimes e) &= T'(f) \\ &= \sum_{n=1}^{\infty} \mu_n(f \otimes e) y_n \\ &= \sum_{n=1}^{\infty} \int_{[0, 1]} f d\langle e, \mu_n \rangle y_n \end{aligned}$$

This of course shows that T' is represented by $\sum_n \langle e, \mu_n \rangle \otimes y_n$; moreover

$$\sum_{n=1}^{\infty} \|\langle e, \mu_n \rangle\| \|y_n\| \leq \|e\| \sum_n \|\mu_n\| \|y_n\| < \infty$$

which implies that T' is nuclear. This contradiction finishes the proof.

This brings us to ask the following question. Let E and F be Banach spaces such that F has the Radon-Nikodym property. Let Ω be a compact Hausdorff space and let

$T : C(\Omega, E) \rightarrow F$ be a bounded linear operator satisfying conditions (i) and (ii) of Proposition 2. Is T nuclear? Recently, the first named author has given a positive answer to the above question when F is complemented in its bidual F^{**} (see [4]).

2–The Radon-Nikodym Property and Nuclear Operators

Throughout this section Ω is a compact Hausdorff space and E and F are Banach spaces. Every bounded linear operator $T : C(\Omega, E) \rightarrow F$ induces a bounded linear operator $T^\# : C(\Omega) \rightarrow \mathcal{L}(E, F)$ where for each f in $C(\Omega)$

$$T^\#(f)(e) = T(f \otimes e)$$

for all $e \in E$.

In this section we shall look at the interplay of the two operators T and $T^\#$. The next result shows that when T is nuclear, then the range of $T^\#$ is in $\mathcal{N}(E, F)$.

Theorem 3 If $T : C(\Omega, E) \rightarrow F$ is nuclear, then $T^\#$ takes its values in $\mathcal{N}(E, F)$.

Proof Let (μ_n) in $M(\Omega, E^*)$ and (y_n) in F be such that $\sum_{n=1}^{\infty} \|\mu_n\| \|y_n\| < \infty$ and for each φ in $C(\Omega, E)$

$$T(\varphi) = \sum_{n=1}^{\infty} \mu_n(\varphi) y_n.$$

For each $n \geq 1$ define $\mu_n^\# : C(\Omega) \rightarrow E^*$ by

$$\mu_n^\#(f)e = \mu_n(f \otimes e)$$

for each $f \in C(\Omega)$ and $e \in E$. Since for each $f \in C(\Omega)$ and $e \in E$

$$\begin{aligned} T^\#(f)(e) &= T(f \otimes e) \\ &= \sum_{n=1}^{\infty} \mu_n(f \otimes e) y_n \\ &= \sum_{n=1}^{\infty} \mu_n^\#(f)(e) y_n \end{aligned}$$

it follows that for each f in $C(\Omega)$ $T^\#(f)$ can be represented by the series $\sum_{n=1}^{\infty} \mu_n^\#(f) \otimes y_n$. Moreover since $\sum_{n=1}^{\infty} \|\mu_n^\#(f)\| \|y_n\| \leq \|f\| \sum_{n=1}^{\infty} \|\mu_n\| \|f_n\| < \infty$, it follows that $T^\#(f)$ is a nuclear operator from E to F for each $f \in C(\Omega)$.

The next result illustrates one key relationship between T and $T^\#$.

Theorem 4 The operator $T : C(\Omega, E) \rightarrow F$ is nuclear whenever $T^\# : C(\Omega) \rightarrow \mathcal{N}(E, F)$ is nuclear.

Proof: Assume $T^\# : C(\Omega) \rightarrow \mathcal{N}(E, F)$ is nuclear. Then there exist sequences (ν_n) in $C(\Omega)^*$ and (N_n) in $\mathcal{N}(E, F)$ such that for each $f \in C(\Omega)$

$$T^\#(f) = \sum_{n=1}^{\infty} \nu_n(f) N_n$$

and

$$\sum_{n=1}^{\infty} \|\nu_n\| \|N_n\|_{\text{nuc}} < \infty.$$

Without loss of generality we may and do assume that $\|\nu_n\| \leq 1$ for all $n \geq 1$. Similarly, since each N_n is a nuclear operator, for each $n \geq 1$, there are sequences $(e_{n,m}^*)$ in E^* and $(y_{n,m})$ in F such that for all $e \in E$

$$N_n(e) = \sum_{m=1}^{\infty} e_{n,m}^*(e) y_{n,m}$$

and

$$\sum_{m=1}^{\infty} \|e_{n,m}^*\| \|y_{n,m}\| \leq \|N_n\|_{\text{nuc}} + 1/2^n.$$

Hence

$$(\dagger\dagger) \quad \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \|\nu_n\| \|e_{n,m}^*\| \|y_{n,m}\| \leq \sum_{n=1}^{\infty} \|\nu_n\| \|N_n\|_{\text{nuc}} + \sum_{n=1}^{\infty} \|\nu_n\| / 2^n < \infty$$

Moreover, for each $f \in C(\Omega)$ and $e \in E$

$$\begin{aligned}
T(f \otimes e) &= T^\#(f)(e) \\
&= \left(\sum_{n=1}^{\infty} \nu_n(f) N_n \right) (e) \\
&= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \nu_n(f) e_{n,m}^*(e) y_{n,m} \\
&= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \nu_n \otimes e_{n,m}^*(f \otimes e) y_{n,m}
\end{aligned}$$

Since the set $\{f \otimes e : f \in C(\Omega), e \in E\}$ is total in $C(\Omega, E)$, we can assert that T is represented by the double indexed series $\sum_n \sum_m (\nu_n \otimes e_{n,m}^*) \otimes y_{n,m}$, where $(\nu_n \otimes e_{n,m}^*)$ is in $M(\Omega, E^*) \simeq C(\Omega, E)^*$ and $(y_{n,m})$ is in F . An appeal to $(\dagger\dagger)$ shows that T is nuclear.

In the following we will show that the converse of Theorem 4 does not always hold. But if in addition E^* is assumed to have the Radon-Nikodym property then a close look at [1, Theorem III.4] reveals that any nuclear operator $T : C(\Omega, E) \rightarrow F$ will indeed induce a nuclear operator $T^\# : C(\Omega) \rightarrow \mathcal{N}(E, F)$. Moreover our next result shows how critical the condition on E^* to have the Radon-Nikodym property in order that T nuclear implies $T^\#$ nuclear. As a matter of fact one can characterize dual Banach spaces with the Radon-Nikodym property as follows:

Theorem 5 Let E and F be Banach spaces. The following properties are equivalent:

- (i) The dual space E^* has (RNP);
- (ii) For every compact space Ω , a bounded linear operator $T : C(\Omega, E) \rightarrow F$ is nuclear if and only if $T^\# : C(\Omega) \rightarrow \mathcal{N}(E, F)$ is nuclear.

Proof: (ii) $\Rightarrow$ (i).

If E^* fails RNP, by [3, p.175] there exists $\mu \in M([0, 1], E^*)$ such that the operator $\mu^\# : C[0, 1] \rightarrow E^*$ defined by

$$\mu^\#(g)e = \mu(g \otimes e)$$

for all $g \in C[0, 1]$ and $e \in E$, is not nuclear. Choose $y \in F$ such that $y \neq 0$, and define $T : C([0, 1], E) \rightarrow F$ by

$$T(\varphi) = \mu(\varphi)y$$

for all $\varphi \in C([0, 1], E)$. It is clear that T is a rank-one operator, hence it is nuclear. The operator $T^\# : C[0, 1] \rightarrow \mathcal{N}(E, F)$ induced by T is clearly the operator such that for each $g \in C[0, 1]$

$$T^\#(g) = \mu^\#(g) \otimes y.$$

To see that $T^\#$ is not nuclear, note that for each $y^* \in F^*$, we can define the operator $T_{y^*}^\# : C[0, 1] \rightarrow E^*$ by

$$T_{y^*}^\#(g) = (T^\#g)^*(y^*)$$

for each $g \in C[0, 1]$.

If $T^\#$ were a nuclear operator then $T_{y^*}^\#$ would also be a nuclear operator for each $y^* \in F^*$. This of course follows from the fact that $T_{y^*}^\#$ is the composition of $T^\#$ and the bounded linear operator from $\mathcal{N}(E, F)$ to E^* which to an element N in $\mathcal{N}(E, F)$ associates the element $N^*(y^*)$ in E^* . But for each g in $C[0, 1]$

$$T_{y^*}^\#(g) = y^*(y)\mu^\#(g).$$

By the Hahn-Banach Theorem, choose y^* in F^* such that $y^*(y) = 1$, then for this particular y^* we have

$$T_{y^*}^\# = \mu^\#.$$

This contradiction shows that $T^\#$ can not be nuclear. This proves (ii) $\Rightarrow$ (i).

The proof of (i) $\Rightarrow$ (ii) is implicit in [1, Theorem III.4]. We shall provide a sketch of a proof for the sake of completeness. For this, assume that $T : C(\Omega, E) \rightarrow F$ is nuclear and that E^* has the Radon-Nikodym property. By Proposition 1 we know that the measure G representing the operator T is countably additive as a vector measure taking its values in $\mathcal{N}(E, F)$ and $|G|_{\text{nuc}}(\Omega) < \infty$. Here it is easy to note that it follows from general vector

measure techniques [3, p.3] that $|G|_{\text{nuc}}$ is countably additive. The proof that $T^\#$ is nuclear now follows the proof of the scalar case as given in [3, p.173] with some minor changes. For instance, since E^* has the Radon-Nikodym property, one can proceed to produce a Bochner $|G|_{\text{nuc}}$ -integrable function

$$H : \Omega \longrightarrow \mathcal{N}(E, F)$$

such that for each Borel subset B of Ω ,

$$G(B) = \int_B H(\omega) d|G|_{\text{nuc}}(\omega).$$

and for each $f \in C(\Omega)$ and each $e \in E$

$$T(f \otimes e) = \int_\Omega f(\omega) H(\omega)(e) d|G|_{\text{nuc}}(\omega).$$

Hence for each $f \in C(\Omega)$

$$T^\#(f) = \int_\Omega f(\omega) H(\omega) d|G|_{\text{nuc}}(\omega)$$

Another appeal to [3, p.173] shows that $T^\#$ is nuclear.

Bibliography

- [1] G. Alexander, *Linear Operators on the Spaces of Vector-Valued Continuous Functions*, Ph. Dissertation, New Mexico State University, Las Cruces, New Mexico, 1972.
- [2] R. Bilyeu and P. Lewis, *Some mapping properties of representing measures*, Ann. Mat. Pure. Appl. CIX (1976) pp. 273–287.
- [3] J. Diestel, and J.J. Uhl, Jr. **Vector Measures**, Math Surveys, **15**, AMS, Providence, RI (1977).
- [4] P. Saab, *Integral Operators on Spaces of Continuous Vector-Valued Functions*, Proc. Amer. Math. Soc. (to appear).
- [5] L. Schwartz, *Séminaire Schwartz, Exposé 13*, Université de Paris, Faculté des Sciences, (1953/54).

University of Missouri
Mathematics Department
Columbia, MO 65211